



Expression of miR-15a-5p, miR-17-5p, and miR-103-3p in hypertension-associated chronic kidney disease and hemodialysis patients

Heba Nazeir¹, Amr E. Ahmed¹, Nahla S. Kotb², Doaa Ezzat³, Mohamed Gamal⁴, Mohamed D.E. Abdelmaksoud⁵, Weaam Gouda^{5*}

¹Department of Biotechnology and Life Sciences, Faculty of Postgraduate Studies for Advanced Sciences, Beni-Suef University, Egypt.

²Department of Biochemistry, Egyptian Drug Authority, Giza, Egypt.

³Medical Biochemistry and Molecular Biology Department, College of Oral and Dental Surgery, Misr University for Science and technology, Egypt.

⁴Internal Medicine Gastroenterology and Hepatology Department, Faculty of Medicine, Ain Shams University, Cairo, Egypt.

⁵Biochemistry Department, Biotechnology Research Institute, National Research Centre, Giza, Egypt.

Corresponding Author:

Name : Weaam Gouda

Address: National Research Centre, Giza, Egypt

Received: 7 February 2025

Revised: 17 March 2025

Accepted: 17 March 2025

Published: 31 August 2025

Egyptian Pharmaceutical Journal 2025, 24: 62-74

Background

MicroRNAs (miRs), which can be released into the extracellular environment, are important modulators of how cells respond to various stimuli. Prior research over the past decade has focused on miRs as new clinical targets or diagnostic tools. Numerous studies on chronic kidney disease (CKD) have focused on miRs. One prevalent risk factor for CKD is hypertension (HTN), which damages targeted organs, including the kidneys, heart, and arteries.

Objective

Our goal was to investigate the expressions of miRs (17-5p, 15a-5p, and 103-3p) in prerenal failure and renal failure patients under hypertension conditions, as these could be biomarkers of kidney damage caused by hypertension.

Patients and methods

Quantitative real-time polymerase chain reaction (RT-PCR) was utilized to assess the expression levels of the above-mentioned miRs. A total of 110 studied cases that met the eligibility criteria of HTN-associated CKD were grouped into a pre-renal failure group (n = 50) and a renal failure group (n = 60), which included end-stage renal disease cases under hemodialysis beside 40 healthy controls.

Results and conclusion

Our research revealed that miR-103-3p was upregulated; however, miRs (15a-5p and 17-5p) were significantly downregulated in the studied HTN-CKD groups compared with control subjects. The degree of deregulation correlated with the severity of the disease suggests that these miRs may be involved in the pathophysiology of HTN-related CKD, which in turn encourages additional research into their possible clinical use in the prevention and management of CKD. Moreover, these miRs showed areas under the curve near 1 in receiver operating characteristics (ROC) analysis. Consequently, they might be used as markers for CKD.

Keywords: chronic kidney disease (CKD), hypertension (HTN), microRNAs (miRs), miR-103-3p, miR-15a-5p, miR-17-5p.

Egypt Pharmaceut J 24: 62-74

© 2025 Egyptian Pharmaceutical Journal

1687-4315

Introduction

Chronic kidney disease (CKD) encompasses all levels of loss in kidney function, ranging from mild to severe chronic renal failure [1]. CKD is defined as a progressive decline in renal function that eventually results in kidney impairment or a glomerular filtration rate (GFR) of less than 60 ml/min per 1.73 m² for more than three months[2]. One of the most prevalent primary conditions that leads to CKD and eventually end-stage renal disease (ESRD) is hypertension (HTN)[3]. There is an epidemiological relationship between HTN and CKD. The prevalence of HTN has been reported to

be greater than 85% in stage 3 and above 90% are in stage 4 and stage 5 CKD patients [4]. Most patients with mild to moderate CKD have no obvious clinical symptoms, so it is difficult to diagnose and as the disease progresses further it may develop into ESRD, which may be irreversible[5]. Albuminuria and GFR are the primary criteria used to diagnose and stage CKD; these tools, however, are mostly capable of diagnosing CKD after renal damage has occurred. They merely offer a limited amount of information regarding kidney disease activities and do not offer any etiological clues [6]; therefore, in order to

lower CKD-related mortality, earlier CKD diagnosis and therapy may be more appropriate. Thus, a deeper insight into the mechanisms behind the onset and progression of CKD is required. Recently, many microRNA (miRs) molecules have been discovered that play an essential role in the pathogenesis of many diseases. Investigating miRs as biomarkers for CKD has emerged as a fascinating area of research [7]. MiRs are defined as short non-coding RNAs that bind to the 3' untranslated regions of their target mRNAs, causing translational repression and mRNA degradation. They are important modulators of gene expression [8], which play a very important and extensive role in a number of cellular regulatory processes, including differentiation, proliferation, development, and apoptosis [9]. They are also involved in the development and proper operation of the kidneys, including the maintenance of fluid, electrolyte, acid-base, and blood pressure [10]. In many biological fluids, these RNA species have longer half-lives and are more stable, such as urine, plasma, serum, and saliva [11]. Therefore, miRs dysregulation may result in cell dysfunction and development of CKD [12-15]. Previous investigations also showed changes in miR expression in the onset and progression of hypertensive nephropathy [16-20]. Prior study revealed that the miR-103 expressions were increased in the plasma of hypertension cases relative to that of normal individuals [18]. A previous study conducted in the United States of America revealed dysregulation of miR-15a-5p and miR-17-5p in hypertensive patients with CKD [21]. The role of miRs (103-3p, 15a-5p, and 17-5p) in HTN associated with CKD needs more investigation in different populations. Hence, the objective of this study is to assess the levels of miR-15a-5p, miR-103-3p, and miR-17-5p in Egyptian CKD patients associated HTN and to evaluate the effect of these miRs levels in the progress of ESRD, which should assist shed light on how ESRD develops and establish a relationship with kidney function and HTN.

Subjects and methods

Study population

The EL-Edwa Hospital, Minia, Egypt, accepted the study protocol and each participant gave their informed consent. The subjects were assigned into three groups according to kidney function test and estimated glomerular filtration rate (eGFR) [22]. Group 1 (control group) consisted of forty healthy volunteers with an age range of 45 to 63 years; all of them had no relevant medical history and were not taking any medication on a regular basis, with an eGFR > 90 mL/min, serum creatinine up to 1.3 mg/dL, and normal blood pressure (BP). Group II (pre-renal failure group) included fifty hypertensive patients with chronic kidney disease (CKD), age range 43-62 years, untreated newly diagnosed

hypertension, with $15 < \text{eGFR} < 60$ mL/min, and serum creatinine ranged from 1.3 to 3 mg/dL. Group III (renal failure group): sixty patients (age range 47-65 years) with end-stage renal disease (ESRD) with a long history of chronic hypertension, $\text{eGFR} < 15$ mL/min, and creatinine > 4.0 mg/dL. All had received hemodialysis therapy for at least 6 months, three times a week for four hours per session. Hypertension was defined according to the European Society of Hypertension clinical practice guidelines [23]. Patients with comorbid conditions including liver disease, autoimmune diseases, or malignancies, as well as patients with renal disease due to pathologies other than hypertension, excessive smoking, and relevant medications (anti-hypertensive drugs), were excluded from the study. Treatments with corticosteroids, cytotoxic medicines, or non-steroidal anti-inflammatory drugs at the time of the study were additional exclusion factors.

Sample collection and measurements

Samples of blood were taken under fasting conditions from every participant in the morning. Blood was collected in vacutainer EDTA for plasma isolation plus serum tubes for blood biochemistry determined by routine methods. At room temperature, the blood samples were separated at 3000 rpm for ten minutes. The plasma was gathered and kept at -80°C until analysis. Serum creatinine level, age, and sex were utilized to assess the eGFR. Individuals were diagnosed with CKD if their eGFR was less than 60 mL/min/1.73 m².

Molecular analysis

The quantitative real-time PCR (qPCR) method was used to evaluate the chosen miRs. We selected three miRs (miR-103-3p, miR-15a-5p and miR-17-5p) for investigation according to involvement in HTN-CKD-related pathways as reported in pertinent literature. Total RNA was extracted from the plasma after the samples were thawed at room temperature using the miRNeasy Plasma Advanced Kit (Qiagen Inc., Germantown, Maryland, USA) according to the manufacturer's instructions. The extracted RNA concentrations were measured by reading the absorbance at 260 nm using Thermo Fisher Scientific Inc., Waltham, MA, USA NanoDrop 2000c spectrophotometer [24]. MiRs were reverse-transcribed to complementary DNA using cDNA Synthesis kit (Applied Biosystems, Foster, California, USA). MiRs expressions were determined via specific primers illustrated in Table 1 [25-27] and the cDNA product was used as a template for the detection of miR expression by using PCR master mix (SYBR green) and utilizing an applied biosystems, USA, real-time PCR System 2700. A conserved gene U6 served as the internal standard. Data are shown as fold changes ($2^{-\Delta\Delta\text{Ct}}$) [28].

Table 1 Primers for reverse transcriptase polymerase chain reaction (RT-PCR).

MiRNAs	Primer sequences	References
MiRNA-15a-5p	Forward:5'-ACACTCCAGCTGGGTAGCAG CACATAATGGTTTGT-3' Reverse:5'-CTCAACTGG TGTCGTGGAGTCGGCAATTCAGTTGAGCACAAAC-3'	[25]
MiRNA-17-5p	Forward:5'-TGCGCCAAAGTGCTTACAGTGCA-3' Reverse:5'- CCAGTGCAGGGTCCGAGGTATT-3'	[26]
MiRNA-103-3p	Forward:5'-ACACTCCAGCTGGGAGCAGCATTGTAC-3' Reverse:5'-TGGTGTCGTGGAGTCG-3'	[27]

Statistical analysis

All measured variables' distributions were checked for normality. One-way analysis of variance and the student's test were used for continuous values with a normal distribution, and the variables were displayed as mean $\pm$ standard deviation (SD). Pearson's correlation analysis was performed to assess correlation coefficients (*r*). Furthermore, the specificity and sensitivity of the regression models that included the related miRs were evaluated using receiver operating characteristics (ROC). SPSS version 27.0 (SPSSInc, Chicago, IL, USA) was used for all tests. A *p*-value of less than 0.05 was regarded as significant.

Results

The clinical and demographical features of the studied subjects are summarized in Table 2. The participant's ages ranged from 43 to 65 years. There was no significant difference in age (*P* = 0.123) or sex (*P* = 0.67) between control, prerenal failure (PRF), and renal failure (RF) groups. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) were significantly elevated in CKD patients than controls. Likewise, serum creatinine and urea were the highest in RF cases (for creatinine: RF cases = 8.30 ± 2.57 mg/dL; PRF cases = 2.01 ± 0.42 mg/dL; controls = 0.80 ± 0.12 mg/dL, *p* < 0.001) (for urea: RF cases = 144.2 ± 32.99 mg/dL; PRF cases = 77.66 ± 26.85 mg/dL; controls = 23.63 ± 3.53 mg/dL, *p* < 0.001). As expected, the mean eGFR was significantly lower in the RF group (7.53 ± 2.52 ml/min/1.73 m²) than PRF cases (38.14 ± 10.78 ml/min/1.73 m²) and control groups (105.78 ± 16.93 ml/min/1.73 m²) (*p* < 0.001). Similarly, significant differences were observed between the studied groups with respect to hemoglobin (Hb) (*P* = 0.01), and the level of electrolytes was relatively similar across the groups. As shown in Tables 3 & 4 and Figures 1 & 2, the plasma of hypertensive patients in the study

groups contains higher levels of miR-103-3p than those found in healthy individuals; its expression in PRF and RF cases increased 1.397-fold and 2.831-fold, respectively, compared to controls (*p* < 0.001). Alternatively, miR-15a-5p and miR-17-5p are characterized by lower expression levels in CKD cases. There was a 0.722-fold and 1.412-fold decrease in the expression of miR-17-5p in PRF & RF cases, respectively (*p* < 0.001). We also reported a 0.974-fold lower level of miR-15a-5p expression in the RF group (*p* < 0.001) than the PRF group (log2 fold change = -0.482, *p* < 0.001). Correlations between the studied plasma miRs and clinical parameters in the studied groups were evaluated and represented in Table 5 and Figures 3 & 4. Correlation analysis showed that miR-103-3p was positively linked to SBP, serum creatinine, and urea levels and a corresponding negative correlation with eGFR in PRF&RF groups. In contrast, levels of miR-15a-5p and miR-17-5p were negatively related to SBP, serum creatinine, and urea levels, and a significant positive correlation was obtained with eGFR. No other significant relationship was found among the studied miRs and other clinical parameters. The diagnostic utility of the chosen miRs for the CKD and control groups was then assessed using receiver operating characteristic (ROC) curve analyses. Regarding the PRF group, the area under curves (AUCs) which ranged from 0.783 to 0.898 was displayed in Figure 5 and Table 6. The diagnostic effect of the combined panel of the three miRs was then assessed using the frequency table and ROC curves. The 3-miR panel's ROC curve showed a high degree of diagnostic accuracy. (AUC, 0.978; 95% CI, 0.952–1.00; *p* < 0.001), which was superior to that of most of the single miRs. Concerning the RF group, as can be observed in Figure 6 and Table 7, the AUCs ranged from 0.893 to 0.995, and the ROC curve for the 3-miR panel had the highest AUC (AUC, 1.00; 95% CI, 1.00–1.00; *p* < 0.001). These findings imply that these miRs are powerful diagnostic indicators for CKD patients with hypertension.

Table 2 Demographic, clinic and laboratory characteristics among the studied groups.

Variables	Controls(n=40)	PRF group (n=50)	RF group (n=60)	P value
age (years)	50.15± 3.93	52.04 ± 4.17	55.52± 4.61	0.123
Sex (male/female)	30 /10	38 /12	47 /13	0.67
SBP(mm Hg)	116.91±7.11	142.26±15.34	140.83±16.16	0.01
DBP(mm Hg)	78.11±3.77	91.66±7.56	90.93±8.31	0.01
Creatinine (mg / dL)	0.80±0.12	2.01±0.42	8.30±2.57	<0.001
Urea (mg / dL)	23.63±3.53	77.66±26.85	144.2±32.99	<0.001
eGFR (ml /min / 1.73 m ²)	105.78±16.93	38.14±10.78	7.53±2.52	<0.001
Ca ⁺⁺ (mmol/L)	1.18±0.05	1.17±0.07	1.15±0.07	0.06
Na ⁺ (mmol/L)	141.54±4.12	140.84±7.01	136.37±4.53	0.12
k ⁺ (mmol/L)	4.23±0.41	4.27±0.54	5.12±0.83	0.08
Hb(g/dL)	13.01±0.79	12.55±1.18	9.05±1.97	0.01

Data represented as mean ± standard deviation (SD); PRF: Prerenal failure; RF: Renal failure; SBP: systolic blood pressure; DBP: diastolic blood pressure; eGFR: estimated glomerular filtration rate; Hb: hemoglobin.

Table 3 The relative expression for studied miRs of the population recruited.

Variables	Controls(n=40)		PRF group(n=50)		RF group(n=60)		P value
	M ± SD	95 % CI	M ± SD	95 % CI	M ± SD	95 % CI	
miR-103-3p	1.23±0.75	0.99-1.46	3.22±0.28	2.86-3.59	8.72±0.26	7.87-9.56	<0.01
miR-15a-5p	1.13±0.26	1.05-1.21	0.81±0.33	0.72-0.90	0.58±0.19	0.53-0.62	<0.01
miR-17-5p	1.27±0.44	1.13-1.41	0.77±0.34	0.67-0.87	0.48±0.21	0.42-0.53	<0.01

M ± SD: mean ± standard deviation; 95% CI: 95%; confidence interval; PRF: Prerenal failure; RF: Renal failure.

Table 4 The fold change for the expression levels of prerenal and renal failure groups.

Variables	PRF group (n=50)			RF group (n=60)		
	log2 FC	expression change	P value	log2 FC	expression change	P value
miR- 15a-5p	-0.482	down	<0.001	-0.974	Down	<0.001
miR- 17-5p	-0.722	down	<0.001	-1.412	Down	<0.001
miR- 103-3p	1.397	up	<0.001	2.8314	Up	<0.001

FC; fold change; PRF: Prerenal failure; RF: Renal failure.




Fig. 1. Relative expression of evaluated microRNAs (miRs) in the studied groups: prerenal failure (PRF) and renal failure (RF).




Fig. 2. The fold change for the relative expressions in the studied groups; (A) for prerenal failure (PRF) cases and (B) for renal failure (RF) cases.

Table 5. Correlation analysis between the expression of the three studied plasma miRs and clinical parameters in the studied groups using Pearson's correlation analysis.

variables	Pre renal failure group n= 50						Renal failure group n= 60					
	miR-15a-5p		miR-17-5p		miR-103-3p		miR-15a-5p		miR-17-5p		miR-103-3p	
	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value	<i>r</i>	<i>p</i> -value
Age	-0.174	0.534	-0.238	0.311	0.23	0.153	-0.138	0.198	-0.109	0.334	0.089	0.498
SBP	-0.602	<0.001	-0.611	<0.001	0.505	0.001	-0.539	<0.001	-0.57	<0.001	0.501	0.001
DBP	-0.205	0.152	-0.102	0.211	0.304	0.06	-0.102	0.09	-0.136	0.074	0.208	0.07
Hb	0.328	0.069	0.249	0.121	-0.125	0.09	0.037	0.778	0.046	0.727	-0.112	0.395
Creatinine	-0.593	<0.001	-0.695	<0.001	0.714	<0.001	-0.84	<0.001	-0.868	<0.001	0.526	0.005
Urea	-0.464	0.001	-0.626	<0.001	0.588	<0.001	-0.763	<0.001	-0.819	<0.001	0.643	0.030
eGFR	0.443	0.001	0.551	<0.001	-0.681	<0.001	0.808	<0.001	0.778	<0.001	-0.632	<0.001
Na ⁺	-0.156	0.113	-0.199	0.811	0.272	0.09	-0.107	0.958	-0.135	0.304	0.172	0.586
K ⁺	-0.245	0.128	-0.218	0.177	0.084	0.606	-0.169	0.196	-0.122	0.869	0.151	0.701
Ca ⁺⁺	0.212	0.08	0.268	0.095	-0.29	0.069	0.102	0.437	0.061	0.644	-0.146	0.727

r: correlation coefficient; eGFR: estimated glomerular filtration rate SBP: systolic blood pressure; DBP: diastolic blood pressure.

Table 6. Receiver operating characteristic (ROC) curve analysis of plasma miRs assessed in prerenal failure patients

Variable(s)	AUC	SE	<i>P</i> value	95% CI	
				Lower Bound	Upper Bound
miR-17-5p	0.783	0.051	< 0.001	0.683	0.882
miR-15a-5p	0.842	0.041	< 0.001	0.763	0.922
miR-103-3p	0.898	0.033	< 0.001	0.832	0.963
miR-Panel	0.978	0.013	< 0.001	0.952	1.004

ROC: receiver operating characteristic; AUC: area under curve; SE: standard error; CI: confidence interval.

Table 7 Receiver operating characteristic (ROC) curve analysis of plasma miRs estimated in renal failure group.

Variable(s)	AUC	SE	<i>P</i> value	95% CI	
				Lower Bound	Upper Bound
miR-17-5p	0.971	0.015	< 0.001	0.942	1.001
miR-15a-5p	0.893	0.033	< 0.001	0.828	0.958
miR-103-3p	0.995	0.004	< 0.001	0.987	1.003
miR-Panel	1.00	0.00	< 0.001	1.00	1.00

ROC: Receiver operating characteristic; AUC: area under curve; SE: standard error; CI: confidence interval.




Fig. 3. Significant correlations between three measured plasma microRNAs (miRs) and clinical parameters in prerenal failure (PRF) cases.




Fig. 4. Significant correlations between the plasma microRNAs (miRs) and clinical parameters in renal failure (RF) cases.




Fig. 5. Receiver operating curve (ROC) of the three microRNAs (miRs) investigated in patients with prerenal failure (PRF) compared to controls.




Fig. 6. Receiver operating characteristic (ROC) curve analysis of microRNAs (miRs) in renal failure (RF) cases versus controls.

Discussion

It has been reported that miRNAs (miRs) are crucial molecular modulators. One miR molecule is able to control up to 200 distinct mRNA targets. Therefore, these non-coding RNAs play an important role in the regulation of many physiological and pathological cellular processes [29]. Recently, some studies have reported dysregulated patterns of miRs in chronic kidney disease (CKD) in patients with hypertension (HTN); these miRs could be employed as possible biomarkers and may have a role in the pathophysiology of CKD [30-32]. Findings of this study are that three plasma miRs (miR-15a-5p, miR-103-3p, and miR-17-5p) were significantly dysregulated in hypertensive mild CKD individuals and patients with end-stage renal disease (ESRD). Our data displayed that miR-17-5p was significantly downregulated among CKD cases. This circulating miR belongs to a cluster of miRs located in intron 3 of C13orf25 at chromosome 13 [33]. This cluster plays a distinct role in the differentiation and maturation of immune cells [33]. MiR-15a-5p was also significantly downregulated in the studied cases; in contrast, miR-103-3p showed a significant elevation in cases compared to healthy individuals. The identified miR-15a-5p and miR-103-3p belong to the miR-15/107 group, which is widely expressed in many different mammalian tissues [34]. The biological functions of miRs from this group are involved in cell metabolism and division, angiogenesis, and stress response, assuming that deregulation of these miRs is a hallmark of numerous disease states [35-37]. We found in our study that expression levels of miR-15a-5p and miR-17-5p significantly decreased in ESRD cases than mild CKD individuals. We analyzed the link between eGFR and the expressions of miRs (15a-5p and 17-5p), and we observed that the expression levels of both miRs (15a-5p and 17-5p) decreased as kidney function deteriorated among the studied groups [38]. The decreased level of miRs may be due to their degradation by circulating RNase [39]. In the present study, a negative correlation between SBP and these two miRs was detected in the studied groups. In the same line with our study, previous investigations have shown that expression of these miRs is critical for immune cell function and demonstrated that the expression of miR-17 plays an important role in the production of T cell survival [40,41], while the expression of miR-15 can enhance the induction of regulatory T cells [42]. Likewise, the components of the immune system have a direct impact on the release of inflammatory cytokines and reactive oxygen species in hypertensive conditions [43-47], which are the main contributors to the deterioration of

renal function, which leads to progressive kidney injury [48-50]. The reduction in expression of miRs (15a-5p and 17-5p) in HTN-CKD cases induced immune dysfunction that is well established among end-stage renal disease patients [51]. Thus, downregulation of these miRs in the context of hypertension may lead to the onset of CKD and have consequences for the development of CKD or its complications [21]. Moreover, another study consistent with our finding illustrated that plasma miR-15b levels in hemodialysis patients were 2-fold lower compared to non-CKD, and the author referred to the lower level of miR-15b as being linked to the development of ESRD by modifying phosphate metabolism-related genes [52]. Also, there have been reports of associations between acute kidney injury (AKI) in human and animal models and the levels of miR-17 and miR-15 b within kidney tissues [53]. On contrast, we found in the current study upregulation of miR-103-3p in mild HTN-CKD and ESRD patients; this upregulation was comparable to a study by Lu et al. that showed that, in comparison to healthy controls, hypertensive mice infused with angiotensin II hormone and patients with hypertensive nephropathy had overexpression of miR-103a3p in both serum and urine [18]. Our data also showed a positive correlation of miR-103-3p with SBP. According to our results, some previous studies reported that angiotensin-II levels are elevated in hypertension conditions; this hormone has pro-inflammatory and vasoconstrictive effects on post-glomerular arteries, resulting in kidney fibrosis and glomerular damage [54,55] and this renal injury is characterized by increased renal inflammation, glomerular fibrosis, extracellular matrix deposition, and glomerular protein filtration [56,57]. Lu and colleagues conducted additional studies to investigate the function of miR-103a-3p in angiotensin II-induced renal damage. They found that anti-miR-103a-3p knockdown decreased the effects of angiotensin II on the kidney, while overexpression caused kidney inflammation, kidney fibrosis, and albuminuria in animal models [18]. A target of miR-103a-3p, sucrose non-fermentable-related serine/threonine-protein kinase (SNRK), is expressed less when miR-103a-3p is present. SNRK has the ability to reduce inflammation. Mice with renal damage caused by angiotensin II also had lower SNRK levels; therefore, in contradiction to our finding, Lu et al. discovered that under hypertensive conditions, the levels of angiotensin-II and miR-103 were elevated and the expression of SNRK was decreased compared with healthy kidneys, and these conditions accelerated the progression of CKD [18]. Finally, depending on the estimated cutoff values of the area under the curve (AUC) of the receiver operating characteristic

(ROC) curve, all three miRs had an acceptable ability to predict CKD but miR-103-3p was excellent diagnostic biomarkers for mild HTN-CKD and ESRD cases, which showed a high predictive value however, the prediction model containing three tested miRs offered an additional advantage in predicting CKD in two studied groups. Our results were in consistent with studies by Garmaa *et al* that referred to the fact that miRs can be used as effective diagnostic markers for CKD and the use of a panel of miRs allowed more accurate diagnosis than relying on a single miR [58].

In addition, as all samples in the study were obtained from clinical patients, we suspect that these miRNAs may be more relevant than those that have been identified in animal models, and may therefore be better able to elucidate the putative function of the miRNAs assessed in chronic kidney disease. However, the limitations of the study include the small sample size and the need to further validate the findings in a large patient cohort.

Conclusion

The current study demonstrated that miRNA-103-3p was up-regulated while miRNA-15a-5p and miRNA-17-5p were down-regulated in studied hypertension (HTN) associated chronic kidney disease (CKD) cases. The dysregulation of the mentioned miRNAs was positively associated with prevalent chronic kidney disease in hypertensive individuals and associated with the development of end-stage renal disease (ESRD). Furthermore, we showed that each miRNA expression level was useful for predicting chronic kidney disease (CKD) under hypertension conditions. Thus, these miRNAs could develop into useful non-invasive biomarkers for hypertension (HTN) - chronic kidney disease (CKD); however, more thorough and extensive research is still required to determine the precise connection between miRNAs and hypertension related chronic kidney disease.

Acknowledgements

The National Research Centre's technical support of the current study is gratefully acknowledged. We appreciate the patients and research participants.

Authors' contributions

Concepts and Design: A.A., N.S.K., M.D.E.A., and W.G.; Definition of intellectual content: M.G.; literature search, clinical studies, and data acquisition: H.N., W.G., and D.A.; data analysis and statistical analysis: H.N.; manuscript preparation: H.N. and W.G.; manuscript editing and

review: N.S.K., M.D.E.A., A.A., and W.G.; Guarantor: A.A. and N.S.K.

Conflicts of interest

The authors declare there are no conflicts of interest.

Ethical considerations

This study was performed in line with the principles of the declaration of Helsinki. The study clearance was applied from the Institutional Review Board of the Faculty of Medicine, Minia University (approval No. 213-2022).

References

- Vaidya SR, Aeddula NR. Chronic Kidney Disease. [Updated 2024 Jul 31]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024.
- Chapter 1: definition and classification of CKD. *Kidney IntSuppl* 2013; 3:19–62.
- Webster AC, Nagler EV, Morton RL, Masson P. Chronic Kidney Disease. *Lancet* 2017;389(10075):1238-1252.
- Rao MV, Qiu Y, Wang C, Bakris G. Hypertension and CKD: Kidney Early Evaluation Program (KEEP) and National Health and Nutrition Examination Survey (NHANES), 1999-2004. *Am J Kidney Dis* 2008; 51(4 Suppl 2):S30-7.
- Essue BM, Jha V, John O, Knight J, Jan S. Universal health coverage and chronic kidney disease in India. *Bull. World Health Organ* 2018; 96: 442.
- Al Rubeaan K, Siddiqui K, Al-Ghonaim MA, Youssef AM, Al-Sharqawi AH, AlNaqeb D. Assessment of the diagnostic value of different biomarkers in relation to various stages of diabetic nephropathy in type 2 diabetic patients. *Sci. Rep* 2017; 7: 2684.
- Ata A, Ezzat D, Sherkawy HS, Ahmed AE, Afify M, Abdelmaksoud MDE, *et al.* Impact of micro-RNAs as biomarkers for end-stage renal disease related to hypertension and diabetes. *Sci. Rep* 2025; 15(1): 31101.
- Kim VN, Han J, Siomi MC. Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol* 2009; 10:126–139.
- Bartel DP. MicroRNAs: Target recognition and regulatory functions. *Cell* 2009; 136: 215–233.
- Bhatt K, Mi QS, Dong Z. MicroRNAs in kidneys: Biogenesis, regulation, and pathophysiological roles. *Am. J. Physiol.-Ren. Physiol* 2011; 300: 602–610.
- Keller S, Ridinger J, Rupp AK, Janssen JW, Altevogt P. Body fluid derived exosomes as a novel template for clinical diagnostics. *J. Transl. Med* 2011; 9: 86.
- Muralidharan J, Ramezani A, Hubal M, Knoblach S, Shrivastav S, Karandish S, *et al.* Extracellular microRNA signature in chronic kidney disease. *Am J Physiol Renal Physiol* 2017; 1;312(6): F982-F991
- Rudnicki M, Perco P, D Haene B, Leierer J, Heinzel A, Mühlberger I, *et al.* Renal microRNA- and RNA-profiles in progressive chronic kidney disease. *Eur J Clin Invest* 2016 ;46(3):213-26
- Huttenhofer A, Mayer G. Circulating miRNAs as biomarkers of kidney disease. *Clin. Kidney J* 2017; 10: 27–29.

15. Rysz J, Gluba-Brzozka A, Franczyk B, Jablonowski Z, Cialkowska-Rysz A. Novel biomarkers in the diagnosis of chronic kidney disease and the prediction of its outcome. *Int. J. Mol. Sci* 2017; 18: E1702.
16. Chen C, Lu C, Qian Y, Li H, Tan Y, Cai L, et al. Urinary miR-21 as a potential biomarker of hypertensive kidney injury and fibrosis. *Sci. Rep* 2017; 7: 17737.
17. Lyu H, Li X, Wu Q, Hao L. Overexpression of microRNA-21 mediates Ang II-induced renal fibrosis by activating the TGF-beta1/Smad3 pathway via suppressing PPARalpha. *J. Pharmacol. Sci* 2019; 141: 70–78.
18. Lu Q, Ma Z, Ding Y, Bedarida T, Chen L, Xie Z, et al. Circulating miR-103a-3p contributes to angiotensin II-induced renal inflammation and fibrosis via a SNRK/NF-kappaB/p65 regulatory axis. *Nat. Commun* 2019; 10: 2145.
19. Baker MA, Wang F, Liu Y, Kriegl AJ, Geurts AM, Usa K, et al. MiR-192-5p in the Kidney Protects Against the Development of Hypertension. *Hypertension* 2019; 73: 399–406.
20. Wang Z, Zhu Q, Wang W, Hu J, Li PL, Yi F, et al. Downregulation of microRNA-429 contributes to angiotensin II-induced profibrotic effect in rat kidney. *Am. J. Physiol* 2018; 315: F1536–F1541.
21. Nandakumar P, Tin A, Grove ML, Ma J, Boerwinkle E, Coresh J, et al. MicroRNAs in the miR-17 and miR-15 families are downregulated in chronic kidney disease with hypertension. *PLoS One* 2017; 12(8): e0176734.
22. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int* 2024; 105(4S): S117–S314.
23. Kreutz R, Brunström M, Burnier M, Grassi G, Januszewicz A, Muiesan ML, et al. 2024 European Society of Hypertension clinical practice guidelines for the management of arterial hypertension. *Eur J Intern Med* 2024; 126: 1–15.
24. Hassan R, Husin A, Sulong S, Yusoff S, Johan MF, Yahaya BH, et al. Department of Standard Malaysia, MOSTI. Guidelines for nucleic acid detection and analysis in hematological disorders. *Malays J Pathol* 2015; 37(2): 165–73.
25. Li Y, Liu S, Zhang F, Jiang P, Wu X, Liang Y. Expression of the microRNAs hsa-miR-15a and hsa-miR-16-1 in lens epithelial cells of patients with age-related cataract. *Int J Clin Exp Med* 2015; 8(2): 2405–10.
26. Song J, Liu Y, Wang T, Li B, Zhang S. MiR-17-5p promotes cellular proliferation and invasiveness by targeting RUNX3 in gastric cancer. *Biomed Pharmacother* 2020; 128: 110246.
27. Ding J, Xia C, Cen P, Li S, Yu L, Zhu J, et al. MiR-103-3p promotes hepatic steatosis to aggravate nonalcoholic fatty liver disease by targeting of ACOX1. *Mol Biol Rep* 2022; 49(8): 7297–7305.
28. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001; 25(4): 402–8.
29. Pogribny IP. MicroRNAs as biomarkers for clinical studies. *Exp. Biol. Med* 2018; 243: 283–290.
30. Liu Y, Usa K, Wang F, Liu P, Geurts AM, Li J, et al. MicroRNA-214-3p in the Kidney Contributes to the Development of Hypertension. *J Am Soc Nephrol* 2018; 29(10): 2518–2528.
31. Cheng Y, Wang D, Wang F, Liu J, Huang B, Baker MA, et al. Endogenous miR-204 protects the kidney against chronic injury in hypertension and diabetes. *Am. Soc. Nephrol* 2020; 31: 1539–1554.
32. Berillo O, Huo KG, Fraulob-Aquino JC, Richer C, Briet M, Boutouyrie P, et al. Circulating let-7g-5p and miR-191-5p are independent predictors of chronic kidney disease in hypertensive patients. *Am. J. Hypertens* 2020; 33: 505–513.
33. Mogilyansky E, Rigoutsos I. The miR-17/92 cluster: a comprehensive update on its genomics, genetics, functions and increasingly important and numerous roles in health and disease. *Cell Death Differ* 2013; 20(12): 1603–14.
34. Finnerty JR, Wang WX, Hébert SS, Wilfred BR, Mao G, Nelson PT. The miR-15/107 group of microRNA genes: evolutionary biology, cellular functions, and roles in human diseases. *J Mol Biol* 2010; 402(3): 491–509.
35. Aqeilan RI, Calin GA, Croce CM. miR-15a and miR-16-1 in cancer: discovery, function and future perspectives. *Cell Death Differ* 2010; 17(2): 215–20.
36. Li D, Zhao Y, Liu C, Chen X, Qi Y, Jiang Y, et al. Analysis of MiR-195 and MiR-497 expression, regulation and role in breast cancer. *Clin Cancer Res* 2011; 17(7): 1722–30.
37. Furuta M, Kozaki K, Tanimoto K, Tanaka S, Arai S, Shimamura T, et al. The tumor-suppressive miR-497-195 cluster targets multiple cell-cycle regulators in hepatocellular carcinoma. *PLoS One* 2013; 8(3): e60155.
38. Neal CS, Michael MZ, Pimlott LK, Yong TY, Li JY, Gleadow JM. Circulating microRNA expression is reduced in chronic kidney disease. *Nephrol Dial Transplant* 2011; 26(11): 3794–802.
39. Metzinger-Le V, MBaya-Moutoula E, Taibi F, Massy Z, Metzinger L. Implication of MicroRNAs in the Pathophysiology of Cardiac and Vascular Smooth Muscle Cells. In: *Tech 2012; Current Basic and Pathological Approaches to the Function of Muscle Cells and Tissues - From Molecules to Humans*, chapter 9.
40. Jiang S, Li C, Olive V, Lykken E, Feng F, Sevilla J, et al. Molecular dissection of the miR-17-92 cluster's critical dual roles in promoting Th1 responses and preventing inducible Treg differentiation. *Blood* 2011; 118(20): 5487–97.
41. de Kouchkovsky D, Esensten JH, Rosenthal WL, Morar MM, Bluestone JA, Jeker LT. microRNA-17-92 regulates IL-10 production by regulatory T cells and control of experimental autoimmune encephalomyelitis. *J Immunol Baltim Md* 1950. 2013; 191(4): 1594–605.
42. Singh Y, Garden OA, Lang F, Cobb BS. MicroRNA-15b/16 Enhances the Induction of Regulatory T Cells by Regulating the Expression of Rictor and mTOR. *J Immunol Baltim Md* 1950. 2015; 195(12): 5667–77.
43. Tsukimori K, Nakano H, Wake N. Difference in neutrophil superoxide generation during pregnancy between preeclampsia and essential hypertension. *Hypertension* 2007; 49: 1436–1441.
44. Liao TD, Yang XP, Liu YH, Shesely EG, Cava sin MA, Kuziel WA et al. Role of inflammation in the development of renal damage and dysfunction in

- angiotensin II-induced hypertension. *Hypertension* 2008;52:256–263.
45. Huang L, Wang A, Hao Y, Li W, Liu C, Yang Z, et al. Macrophage Depletion Lowered Blood Pressure and Attenuated Hypertensive Renal Injury and Fibrosis. *Front Physiol* 2018; 7;9:473.
46. Kirabo A, Fontana V, de Faria AP, Loperena R, Galindo CL, Wu J, et al. DC isoketal-modified proteins activate T cells and promote hypertension. *J Clin Invest* 2014;124:4642–4656.
47. Wenzel P, Knorr M, Kossmann S, Stratmann J, Hausding M, Schuhmacher S, et al. Lysozyme M-positive monocytes mediate angiotensin II-induced arterial hypertension and vascular dysfunction. *Circulation* 2011;124:1370–1381
48. Irazabal MV, Torres VE. Reactive Oxygen Species and Redox Signaling in Chronic Kidney Disease. *Cells* 2020;9(6):1342.
49. Verma S, Singh P, Khurana S, Ganguly NK, Kukreti R, Saso L, et al. Implications of oxidative stress in chronic kidney disease: a review on current concepts and therapies. *Kidney Res ClinPract* 2021;40(2):183-193.
50. Agharazii M, St-Louis R, Gautier-Bastien A, Ung RV, Mokas S, Larivière R, et al. Inflammatory cytokines and reactive oxygen species as mediators of chronic kidney disease-related vascular calcification. *Am J Hypertens* 2015 ;28(6):746-55.
51. Chonchol M. Neutrophil dysfunction and infection risk in end stage renal disease. *Semin Dial* 2006;19(4):2916.
52. Wang H, Peng W, Ouyang X, Dai Y. Reduced circulating miR-15b is correlated with phosphate metabolism in patients with end-stage renal disease on maintenance hemodialysis. *Ren Fail* 2012;34(6):685-90
53. Fan PC, Chen CC, Chen YC, Chang YS, Chu PH. MicroRNAs in acute kidney injury. *Hum Genomics* 2016;10: 29.
54. Murphy AM, Wong AL, Bezuhly M. Modulation of angiotensin II signaling in the prevention of fibrosis. *Fibrogenesis Tissue Repair* 2015;8:7.
55. Mezzano SA, Ruiz-Ortega M, Egido J. Angiotensin II and renal fibrosis. *Hypertension* 2001;38(3 Pt 2):635-8.
56. Kitching AR, Hutton H L. The players: cells involved in glomerular disease. *Clin. J. Am. Soc. Nephrol* 2016;11: 1664–1674.
57. Kurts C, Panzer U, Anders HJ, Rees AJ. The immune system and kidney disease: basic concepts and clinical implications. *Nat Rev Immunol* 2013;13(10):738-53.
58. Garmaa G, Nagy R, Kóti T, To UND, Gergő D, Kleiner D, et al. Panel miRNAs are potential diagnostic markers for chronic kidney diseases: a systematic review and meta-analysis. *BMC Nephrol* 2024;13;25(1):261.